



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Occurrence of *Mycobacterium avium* subsp. *paratuberculosis* in fecal samples of hunted deer, chamois and ibex in Switzerland

Sarno, E ; Keller, S ; Wittenbrink, M M ; Stephan, Roger

DOI: <https://doi.org/10.1024/0036-7281/a000504>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-87566>

Journal Article

Accepted Version

Originally published at:

Sarno, E; Keller, S; Wittenbrink, M M; Stephan, Roger (2013). Occurrence of *Mycobacterium avium* subsp. *paratuberculosis* in fecal samples of hunted deer, chamois and ibex in Switzerland. *Schweizer Archiv für Tierheilkunde*, 155:523-525.

DOI: <https://doi.org/10.1024/0036-7281/a000504>

Occurrence of *Mycobacterium avium* subsp. *paratuberculosis* in fecal samples of hunted deer,
chamois and ibex in Switzerland

E. Sarno^{1,2}, S. Keller³, M.M. Wittenbrink³, R. Stephan¹

¹ Institute of Food Safety and Hygiene, University of Zurich, ²Department of Zootechnical
Science and Food Inspection, University of Naples Federico II, Naples, Italy, ³Institute of
Veterinary Bacteriology, University of Zurich

Deutscher Titel für Jahresverzeichnis:

Vorkommen von *Mycobacterium avium* subsp. *paratuberculosis* in Kotproben von gejagten
Rehen, Hirschen, Steinböcken und Gämsen in der Schweiz

23

24 **Introduction**

25 *Mycobacterium avium* subsp. *paratuberculosis* (MAP) is the etiological agent of bovine
26 paratuberculosis, a chronic granulomatous enteritis, also known as Johne's disease. MAP
27 prevalence in domestic food producing animals is increasing globally, causing significant
28 economic losses to the livestock industries through subclinical effects (reduced milk yield and
29 weight loss) and subsequent death of the affected animals. Moreover, several reports suggest
30 a possible link between human Crohn's disease and MAP due to remarkable clinical,
31 epidemiological and pathological similarities (Chacon *et al.*, 2004), but a causal relationship
32 has not yet been proven. Nevertheless, the possible involvement in human disease has led to
33 an increased awareness of these microorganisms as public health concerns.

34 The natural hosts for MAP are domestic ruminants with higher prevalence in dairy and beef
35 cattle herds around the world (Manning and Collins, 2001). For Switzerland, different studies
36 showed a low prevalence of MAP in the cattle population (Glanemann *et al.*, 2004; Bosshard
37 *et al.*, 2006).

38 Wild ruminants may act as a reservoir and contribute to the spread of MAP through fecal
39 shedding (Manning, 2011). Their interaction with susceptible farmed ruminants raises the
40 possibility of playing a role in the epidemiology of the disease in domestic livestock. An
41 interspecies transmission between cattle and wild-living red deer (*Cervus elaphus*) has been
42 recently proposed by multitarget genotyping of strains from both animal species (Fritsch *et al.*,
43 2012). However, reported detection rates in wild ruminants vary among different surveys.
44 From fecal samples of red deer a prevalence of 0.1% has been reported in a Czech Republic
45 survey (Pribylova *et al.*, 2011), whereas a much higher prevalence was found in India where
46 15 % of hog deer (*Axis porcinus*) and 57 % of gaur fecal samples (*Bos gaurus*) were tested
47 positive for MAP (Singh *et al.*, 2010).

In view of detection techniques, several culture methods without decontamination or combining chemical or antibiotic decontamination steps and selective media have been proposed (Glanemann *et al.*, 2004; Akineden *et al.*, 2011). However, these methods are time consuming. Immunological-based systems are faster than culture methods but have a low sensitivity (Glanemann *et al.*, 2004; Collins *et al.*, 2005). Molecular based-methods such as PCR assays may represent a rapid screening system for MAP detection. Most of these protocols are based on the IS900 insertion elements but IS900-like sequences found in other Mycobacteria can lead to false positive results. Therefore other sequences without homologues in other Mycobacteria have been proposed; ISMap02 is present in six copies (Stabel and Bannantine, 2005), ISMap2 is present in three copies (Li *et al.*, 2005), and F57 and Hsp X both occur as single copy genes (Tasara and Stephan, 2005). Nevertheless, these methods have a lower sensitivity compared to the cultural based methods and may not detect subclinically infected animals (low shedders). The objectives of this study were a) to investigate the occurrence of MAP in fecal samples collected from hunted wild ruminants by a combination of cultural detection and PCR based identification and b) to assess the impact of wild ruminants in the spread of MAP in Switzerland.

Sampling

During September to November 2011, state gamekeepers and hunters collected immediately after shooting and evisceration, *in loco*, fecal samples from wild ruminants. For each animal, sex, age, and location of hunting were recorded. Fecal matter (approximately 10 grams per sample) was collected from the colon after the large intestine had been opened, placed into sterile tubes and sent to the laboratory where samples were stored at -20°C until processing.

Detection of MAP

For MAP detection, fecal samples from 3 animals of the same species were pooled (total weight of 6g). Each pooled sample was first homogenized in 8ml of sterile demineralized water and stored under chilled conditions for 4 h. Thereafter, samples were decontaminated using a NaOH/oxalic acid treatment (Beerwerth, 1967). Briefly, samples were mixed in 50 ml of a 4% NaOH solution and subsequently transferred into a sterile centrifugation tube and shaken for 8 min on an orbital shaker (Bühler, Hechingen, Germany). After centrifugation (5010 rpm, 15 min, 20°C) the pellet was resuspended in 20 ml of a 5% oxalic acid solution and shaken for 15 min on an orbital shaker. Tubes were centrifuged a second time as described before; the pellet was resuspended in 6ml of sterile 0.9% saline and used as inoculum. For each sample Herrold's Egg Yolk Agar with Mycobactin J and ANV (HEYA) (Becton, Dickinson and Company, Sparks, MD, USA) in duplicates were inoculated with an aliquot (200 µl) of the final processed and decontaminated material. Slants were incubated at 37°C and monitored visually for growth of presumptive positive MAP colonies (small whitish colonies slightly elevated from the surface) every two weeks for up to 16 weeks. Presumptive positive colonies were recorded, subcultured in Middlebrook 7H9 broth supplemented with Middlebrook-ADC growth supplement (Becton, Dickinson and Company, USA) and 2 mg/L Mycobactin J (Synbiotics Europe SAS, Munich, Germany), and incubated at 37°C for 1 week. After Ziehl-Neelsen staining, subcultures with red acid-fast rods were finally confirmed as MAP by using an F57 sequence-based Real-Time PCR protocol (Bosshard *et al.*, 2006).

Results and Discussion

In total 198 fecal samples (69 red deer (*Cervus elaphus*), 51 roe deer (*Capreolus capreolus*), 51 chamois (*Rupicapra rupicapra*), and 9 ibex (*Capra ibex*)) were collected. The majority of the hunted animals were on average 2 years old and originated from the central and eastern part of Switzerland. Using the described methods, none of the pooled fecal samples from the wild ruminants were found to be positive for viable MAP (Tabl 1).

Detection was based on phenotypic properties (small whitish colony morphology and slow growth) and Mycobactin J dependency. Culture-positive HEYA slants (MAP-like colonies) were found in 19 out of 66 decontaminated fecal pools (11 red deer, 7 roe deer and 1 chamois). None of the ibex samples showed bacterial growth. Red acid fast rods forming cluster from culture-positive HEYA slants were visually detected by Ziehl-Neelsen staining from 10 red deer, 4 roe deer and 1 chamois, respectively, but finally confirmed as non-MAP cells by F57 PCR.

Even though competitive growing microorganisms and chemical treatments might hinder and reduce growth of small numbers of viable MAP cells (Grant *et al.*, 2003), the cultural technique is still considered the “reference standard” for detection of MAP in fecal samples. Based on the results of this first study, a low occurrence of MAP in wild ruminants in Switzerland can currently be postulated. Therefore, wild ruminants do not seem to have a big impact in spreading MAP to farmed ruminants. Nevertheless, further studies covering all regions of Switzerland and including a higher number of animals are necessary to get a complete picture of the impact of wild ruminants in the epidemiology of MAP.

Acknowledgements

We would like to thank state gamekeepers and hunters for providing fecal samples from animals.

References

Akineden Ö., Fernández-Silva J. A., Weirich S., Abdulmawjood A., Bülte M.: Comparison of two decontamination procedures, three culture media, and real time-PCR assay for the detection of *Mycobacterium avium* subsp. *paratuberculosis* (MAP) from artificially contaminated raw sausage. Arch. Lebensmittelhyg. 2011, 62: 150-156.

125 *Beerwerth W.*: Die Züchtung von Mykobakterien aus dem Kot der Haustiere und ihre
126 Bedeutung für die Epidemiologie und Bekämpfung der Tuberkulose. *Prax. Pneumol.* 1967,
127 21: 189-202.
128
129 *Bosshard C., Stephan R., Tasara T.*: Application of an F57 sequence-based real-time PCR
130 assay for *Mycobacterium paratuberculosis* detection in bulk tank raw milk and slaughtered
131 healthy dairy cows. *J. Food Protect.* 2006, 69: 1662-1667.
132
133 *Chacon O., Bermudez L. E., Barletta R. G.*: Johne's disease, inflammatory bowel disease, and
134 *Mycobacterium paratuberculosis*. *Annu. Rev. Microbiol.* 2004, 58: 329-363.
135
136 *Collins M. T., Wells S. J., Petrini K. R., Collins J. E., Schultz R. D., Whitlock R. H.*:
137 Evaluation of five antibody detection tests for diagnosis of bovine paratuberculosis. *Clin.*
138 *Diagn. Lab. Immunol.* 2005, 12: 685-692.
139
140 *Fritsch I., Luyven G., Köhler H., Lutz W., Möbius P.*: Suspicion of *Mycobacterium avium*
141 subsp. *paratuberculosis* transmission between cattle and wild-living red deer (*Cervus*
142 *elaphus*) by multitarget genotyping. *Appl. Environ. Microbiol.* 2012, 78: 1132-1139.
143
144 *Glanemann B., Hoelzle L. E., Bögli-Stuber K., Jemmi T., Wittenbrink M. M.*: Detection of
145 *Mycobacterium avium* subspecies *paratuberculosis* in Swiss dairy cattle by culture and
146 serology. *Schweiz. Arch. Tierheilkd.* 2004, 146: 409-15.
147
148 *Grant I. R., Kirk R. B., Hitchings E., Rowe M. T.*: Comparative evaluation of the MGIT and
149 BACTEC culture systems for the recovery of *Mycobacterium avium* subsp. *paratuberculosis*
150 from milk. *J. Appl. Microbiol.* 2003, 95: 196-201.

151

152 *Li L., Bannantine J. P., Zhang Q., Amonsin A., May B. J., Alt D., Banerji N., Kanjilal S.,*

153 *Kapur V.:* The complete genome sequence of *Mycobacterium avium* subspecies

154 *paratuberculosis*. Proc. Nat. Acad. Sci. 2005, 102: 12344-12349.

155

156 *Manning E. J., Collins M. T.:* *Mycobacterium avium* subsp. *paratuberculosis*: pathogen,

157 pathogenesis and diagnosis. Rev. Sci. Tech. 2001, 20: 133-150.

158

159 *Manning E. J.:* Paratuberculosis in captive and free-ranging wildlife. Vet. Clin. North. Am.

160 Food Anim. Pract. 2011, 27: 621-630.

161

162 *Pribylova R., Lamka J., Kopečna M., Trčka I., Moravkova M., Pavlík I.:* Mycobacterial

163 screening of Czech red deer (*Cervus elaphus*) populations in overwintering sites, 2004-2006.

164 J. Wild. Dis., 2011, 47: 780-783.

165

166 *Singh A. V., Singh S. V., Singh P. K., Sohal J. S.:* Genotype diversity in Indian isolates of

167 *Mycobacterium avium* subspecies *paratuberculosis* recovered from domestic and wild

168 ruminants from different agro-climatic regions. Comp. Immunol. Microbiol. Infect. Dis. 2010,

169 33: 127-131.

170

171 *Stabel J. R., Bannantine J. P.:* Development of a nested PCR method targeting a unique

172 multicopy element, ISMap 02, for detection of *Mycobacterium avium* subsp. *paratuberculosis*

173 in fecal samples. J. Clin. Microbiol. 2005, 43: 4744-4750.

174

175 *Tasara T., Stephan R.:* Development of F57 sequence based Real-Time PCR assay for
176 detection of *Mycobacterium avium* subsp. *paratuberculosis* in milk. Appl. Environ. Microbiol.
177 2005, 71: 5957-5968.

178

179 *Tasara T., Hoelzle L. E., Stephan R.:* Development and evaluation of a *Mycobacterium avium*
180 subspecies *paratuberculosis* (MAP) specific multiplex PCR assay. Int. J. Food Microbiol.
181 2005, 104: 279-287.

182

183

184

185 Corresponding author

186 *Roger Stephan, Institute for Food Safety and Hygiene, Vetsuisse Faculty University of Zurich,*
187 *Winterthurerstr. 272, CH-8057 Zurich, Switzerland* E-mail: stephanr@fsafety.uzh.ch.

188

Table 1: Origin and MAP detection results of the 198 fecal samples collected from wild ruminants during the hunting season 2011

Species	No. Samples	No. Pools	No. of pools with growth of presumptive positive colonies	No. of F57 PCR confirmed MAP positive samples
Red deer	69	23	11	0
Roe deer	51	17	7	0
Chamois	51	17	1	0
Ibex	9	9	0	0
Total	198	66	19	0